

Protoplasts Isolation From Leaf Explants Of *Solanum Surattense* Burm. F. A Medicinally Important Plant

Ayodhya Ramulu.Ch, Venkateshwarlu. M, Srilatha. T., Balaraju.P, Ugandhar.T

Abstract— A simple protoplast isolation protocol that was designed to recover totipotent plant protoplasts with relative ease has been described. The key elements of the protocol are, tissue digestion at slightly elevated temperatures and use of protoplast releasing enzymes that are stable and efficient at higher temperatures. Besides enzymes, the protoplast isolation cocktail consisted of an Osmoticum (Mannitol or $MgSO_4$), and a Protectant ($CaCl_2 \cdot 2H_2O$), all dissolved in distilled water. The protocol has ensured reproducibility, higher yields and is gentle on protoplasts as the protoplasts obtained were amenable to cell wall regeneration and cell division. Plant regeneration was demonstrated for *Solanum surattense* Burm.f from protoplasts isolated by this method. Wall regeneration and cell division were obtained in other species. The merits of the protocol are, simple and easy-to-handle procedure, non-requirement of preconditioning of donor plant and explants, incubation without agitation, satisfactory yields, culturability of the protoplasts isolated and applicability of the protocol to a large number of species including mucilage-containing plants

Index Terms— Plant protoplasts; isolation; simple protocol; totipotency.

I. INTRODUCTION

Solanum surattense Burm.f is a medicinal herb belonging to family Solanaceae distributed in arid and semiarid regions of the world, especially in Southeast Asia, Malay, tropical Australia and India. The plant is used as digestive, diuretic and astringent agent, and in bronchial asthma (Govindan *et al.*, 2004). It is also valued for antispasmodic, antitumor, cardiogenic, hypotensive and anaphylactic activity. *S. surattense* produces glycolalkaloids in all parts of the plant body which on hydrolysis and removal of sugar residues yield steroidal alkaloids solanine, solamargine and solasodine. Solasodine is considered as a potential alternative to diosgenin for commercial steroid drug synthesis like progesterone and cortisone (Galanes *et al.*, 1984).

S. surattense is propagated only by seeds, but this method is beset with difficulties such as: (i) The seeds show a low level of germination under normal conditions (ii) the seeds lose their viability on storage and (iii) seed derived progenies are not true-to-type due to crosspollination. Due to over exploitation for high medicinal values and destruction of the habitat, this plant is becoming endan-gered (Khan and

Frost, 2001), hence there is need for *ex situ* conservation through tissue culture technique. Sinha *et al.*, (1979),

The development of protoplast systems has increased the versatility of plants for use in both biochemical and genetic research. They have become indispensable tools in genetic engineering and crop breeding of all the possible starting points for plant genetic manipulation, only protoplasts offer the opportunity to take advantage of all the technologies now available. Since the first successful isolation of protoplasts by Cocking (1960), substantial progress has been made towards improving the technology. Attempts have also been made to isolate protoplasts from several crop species and protoplast-based plant regeneration systems are made available for a great number of species (Maheshwari *et al* 1986). The improvements that have occurred include modification of protoplast isolation procedures (Mei-Lei *et al* 1987), media composition (Kao and Michayluk 1975), preconditioning of protoplast donor tissues (Shahin 1985), utilization of conditioned media or feeder cells (Bellincampi and Morpugo 1987; Kyojuka *et al* 1987; Lee *et al* 1990), and manipulation of culture environment (d'Utra Vaz *et al* 1992).

The success of a protoplast culture system primarily lies with consistent yields of a large population of uniform and highly viable protoplasts. Several protoplast isolation and purification protocols have been published to optimize the yield and reproducibility. They are often procedures of elaborate nature, labour-intensive involving too many explant or protoplast handling steps, and require extended exposure of explant to digestion environment. Further, the efficacy of such protocols or that of enzyme combinations used therein could be limited to a few plant species. These restrictions must be overcome by improvement of the existing conditions and methods. A number of commercial cellulases and pectinases which allow protoplast release are available. By manipulating the source and concentrations of these, protoplasts may be released from most tissues; however, generalizations cannot be made.

The enzymes and techniques used for isolation of protoplasts have a bearing on their subsequent behaviour and development. Methods with too many steps involved often result in the introduction of cell contamination at some stage or the other. Here, we present a simple method in which slightly elevated temperatures and a set of new enzymes that are efficiency at higher temperatures have functioned synergistically to release protoplasts with relative ease in a number of plant species. The enzymes are hitherto not known as being used for protoplast isolation (Sankara Rao and Srikantha 1986). Consistently high yields of viable protoplasts from variety of explants of taxonomically widely separate plants were demonstrated. The cultivability of these protoplasts was examined. The overall efficiently and relative advantages of the method are discussed.

Manuscript received October 04, 2014.

Ayodhya Ramulu.Ch, Regional Institute of Education, Unit IX, Sachiwalaya marg, Bhubaneswar- 751 022

Venkateshwarlu.M, Department of Botany, University College Kakatiya University Warangal 506009

Srilatha. T, Department of Botany, Govt Pingle College Warangal 506009

Balaraju.P, Department of Botany, Govt. Degree & PG College Peddapally - 505 172

Ugandhar.T, Department of Botany, SRR Govt. Arts & Science College Karimnagar - 505001

The protoplast, also known as naked plant cell refers to all the components of plant cell excluding the cell wall, Hanstein introduced the term protoplast in 1880 to designate the living matter enclosed by plant cell membrane. The isolation of protoplasts from plant cells was first achieved by micro surgery on plasmolyzed cells by mechanical method (Klercker, 1892). Protoplast can be isolated from plant tissues (or) cultured cells by enzymatic digestion to remove the cell wall. Besides been useful for cell fusion studies, higher plant protoplasts can also take up. Through the naked plasma membrane, foreign DNA cell organelles, bacteria or virus particles these unique properties of protoplasts, combined with totipotent nature of plant cells, have opened up an entirely area of fundamental and applied search in experimental biology and somatic cell genetic (Gleddie *et al.*, 1986).

Methods and procedure for protoplast isolation from plant tissues have long been known (Keller *et al.*, 1982) Recent advance in the isolation, culture and regeneration of plants from protoplasts of a wide diversity of species have been reported (Gleddie *et al.*, 1989).

Essential in gradient of the technique of genetic modification of plant cells through the protoplast system are

1. Isolation of protoplast
2. Culture of protoplasts to raise whole plant
3. Cell fusion
4. Introduction of foreign genetic material cell organelles into the protoplasts.

A. Source of material

The most convenient and populous source of plant protoplasts is the leaf because it allows the isolation of large number of relatively uniform cells without the necessity of killing the plants. Moreover, the mesophyll cells are loosely arranged, the enzymes have an easy access to the cell wall. When protoplasts are prepared from leaves, the age of the plant and the conditions under which it has grown may be critical. To achieve maximum control on the growth conditions of source plants several workers have used *in vitro* grown shoots (Binding, 1975; Schieder, 1978a; Butenko and Kuchko, 1980). In some species where it is difficult to isolate culturable protoplasts from leaf cells alternative source material of cultured cells have been used. The yield of protoplasts from cultured cells depends on the growth rate and growth phase of cells. Frequently sub cultured suspension cultures, and cells taken from the early log phase are almost suitable (Vasil and Vasil, 1979).

B. Enzyme treatment

The release of protoplasts is very much dependent on the nature and concentrations of enzymes used. The two enzymes regarded essential to isolate protoplasts from plant cells are cellulose & macerozyme. Driselase, having and number of zymolytic activities such as cellulose, pectinase laminarinase and zylanase has proved especially useful for isolating protoplasts from cultured cells. Increase in yield of mulberry proplasts by treatment with chemical substances has been reported earlier (Ohnishi and Kiyama, 1987).

C. *Osmticum*

A fundamental property of isolated protoplasts is their osmotic fragility and hence the need for suitable osmotic stabilizer to the enzyme solution, the protoplast washing medium and protoplast culture medium is necessary. A variety of solutes, ionic and non-ionic, have been tested for adjusting the osmotic pressure of the various solutions used in protoplast isolation and culture, but the most widely used osmotica are sorbitol and mannitol.

With advancement in plant tissue culture technology, plant regeneration systems from protoplasts have been developed in may mulberry species (Ohyama and Oka, 1975). Establishment of protoplast regeneration system using new technologies such as protoplast fusion and gene transfer would contribute to the improvement of mulberry varieties.

In the present study, attempts have seen made to study protoplast isolation and purification using leaf explants of *S. surattense*

II. MATERIAL AND METHODS

A. Plant material

Seeds of *S. surattense* collected from plants growing in the University Campus, Jaipur were used. The seeds were first washed with running tap water, then surface sterilized as follows: (i) Seeds were submerged in 70% (v/v) ethanol (EtOH) for 30 s, (ii) rinsed with sterile distilled water (iii) dipped in sodium hypochlorite (5% w/v) solution for 15 min, finally rinsed with sterile distilled water for three times. The seeds were then germinated on ½ strength Murashige and Skoog (1962) medium containing 1% (w/v) sucrose and 0.8% agar. For shoot regeneration, leaf (1 to 2 cm) excised from 4 week old sterile *in vitro* grown seedling were inoculated onto nutrient medium dispensed in 100 ml Erlenmeyer flasks.

B. Culture media and culture conditions

The pH of MS media used was adjusted to 5.8 using 1 N HCL or 1 N NaOH, before autoclaving at 120°C with 1.5 kgcm-2 pressure for 20 min. The cultures were incubated in a growth chamber at 25 ± 1°C under 16/8 (light/dark) photoperiod with 25 µmol m-2s-1 illumination from cool fluorescent tubes (Philips, India).

The *In vitro* leaves of 2-3 cm in length and 1- 1.5 cm in width were excised from 6 weeks old seedlings. The leaves were cut into pieces smaller than 1mm and incubated in filter sterilized enzyme solution. The enzyme solution consisted of 20% cellulose "onuzuka" R-10 and 1% macerozyme R 10 prepared in MS salts at pH 5.5 with 0.6 m mannitol and osmoticum the sliced leaf pieces of all the cultivars were incubated in 10 ml of enzyme solution at 27°C and shaken at 40- 50 rpm for 4 5 hrs in dark.

C. Isolation of protoplasts from callus cultures

Seeds of *S. surattense* were soaked for 24 hrs in sterilized water and surface sterilized with 0.1% HgCl₂ for 3 to 5 minutes. Then these were washed 3 times with sterile distilled water for 5 minutes and germinated aseptically on MS basal medium. The cotyledon (0.5 – 0.8 cm²) from 4-weeks old axenic – seedlings were excised and inoculated to MS medium supplemented with 2.0 mg/L 2.4.D. alone and gelled with 0.8% agar cultures were maintained at 25 ± 2°C

under a 16 hrs. Photoperiod was provided by cool white fluorescent lamps.

15 – 20 days old. One – gram friable callus derived from cotyledon explants was gently broken into small pieces of callus. Incubation was carried out in 10ml of digestion solution containing 1% cellulose and 0.5% macerozyme with 0.6 m mannitol as osmoticum. Flasks were incubated at 27° C and shaken at 50 rpm for 3-4 hours in dark.

D. Purification of mesophyll and callus derived protoplasts.

The protoplasts were then purified by 60 µ m steel mesh. The filtrate was collected in screw cap centrifuge tube and centrifuged at 50 g for 5 minutes. The supernatant was discarded and the pellet containing protoplasts was loaded on 20% sucrose solution for purification and centrifuged at 100g for 10 minutes to get a distinct protoplast band. The band was taken in a screw cap centrifuge tube and washed with 5ml of 0.6M mannitol by centrifuging at 50g for 5-7 minutes. The pellet was suspended in culture medium containing MS + 0.6 M mannitol + 2, 4-D and BAP at pH 5.7 and the protoplast yield was estimated using hemocytometer. Proto col is adopted for the purification of mesophyll derived protoplasts.

III. RESULTS

A mixture of 2% cellulose and 1% macerozyme was suitable for isolation of viable protoplasts from mesophyll tissues of leaf cultures of *S. surattense* were investigated each enzyme was in effective by itself but when used in combination it resulted satisfactorily (Evans and Bravo 1983). For *in vitro* leaf explants above combination of macerozyme and cellulose gave an optimum yield.




Fig-I Protoplast Isolation in Leaf explants of *S. surattense*: Fresh and viable mesophyll derived Protoplast

In case of *S. surattense* cultivars, prolonged incubation period's i.e 10-12hrs were observed to be unfavorable shrinkage of protoplasts in these cultivars. Cell

digestion was taisy good when above mentioned conditions were applied for protoplast isolation. The number of protoplasts showed increase during shorter treatment time and reached a peak at 4-5 hours of incubation in dark. Beyond 5 hours of incubation the protoplast yield gradually decreased and further resulted in complete shrinkage of protoplasts at 10-12 hours of incubation from these result it was estimate that the adequate time for enzyme treatment to isolate maximum number of protoplasts from mesophyll cells of *S. surattense* (Plate -I).

IV. DISCUSSION

We were successful in protoplast isolation from callus culture and leaf mesophyll cells on the enzyme solution consisted of 2% cellulose "onuzuka" R- 10 and 1 % macerozyme R 10 prepared in MS salts at pH 5.5 with 0.6 M mannitol as osmoticum. Mesophyll cells and callus cells immersed in the enzyme mixture shaken at 40-50 rpm in dark for 10-12 hours resulted in better yield of protoplasts cell digestion was fairly good when above mentioned conditions were applied for protoplast isolation. Several authors have reported that cell digestion and protoplasts yield in case of mulberry was fairly good during 12-13 hours incubation in dark (Tewary and Lakshmisita 1992).

However our findings agree with the results of who have examined a similar effect of enzyme treatment time for shorter duration (10- 12 hours) on protoplast isolation from seedling cotyledons of *S. surattense*. While in the case of callus, cellulose 1% and macerozyme 0.5 % yielded maximum number of protoplast. The use of callus as source for protoplast isolation may overcome the difficulty encountered with the production of leaf material in some recalcitrant *S. surattense* cultivars. Callus developed on 2.0 mg/L 2,4-D was used for protoplast isolation. A part from sub cultured callus the use of cell suspension culture is also avoided here to isolate protoplasts; this is because the suspension culture may accumulate changes in ploidy and abberations due to soma clonal variations (Larkin and Scowcroft 1981).

REFERENCES

- [1] **Bellincampi D and Morpugo G (1987)**. Conditioning factor affecting growth in plant cells in culture; *Plant Sci.* **51** 83-91
- [2] **Binding, H., 1974**): Cell cluster formation by leaf protoplasts from axenic cultures of haploid
- [3] *Petunia hybrida* L. *Plant Sci. Lett.* **2**, 185--188.
- [4] **Butenko R. and Kuchko A.A.** (1980). Physiological aspects of procurement, cultivation and hybridization of isolated potato protoplasts *Sov. Plant Physiol.* (English Transl.), **26**: 901-909.
- [5] **Cocking E C (1960)** A method for the isolation of plant protoplasts and vacuoles; *Nature (London)* **187** 927-929
- [6] **d'Utra Vaz F B, Slamet I H, Khatum A, Cocking E C and Power J B (1992)** Protoplast culture in high molecular oxygen atmosphere; *Plant Cell Rep.* **11** 416-418
- [7] **Evans D.A. and Bravo J.E. (1983)**. Protoplasts, isolation & culture. In: Hand book of plant cell culture, Vol. I Eds., Sharp W.R. Ammiralo P.V. and Yamada Y., *MacMillan Pub. Co., New York*, pp. 124- 176.
- [8] **Galanas IT, Webb DT, Rosario O (1984)**. Steroid production by callus and cell suspension cultures of *Solanum aviculare* Forst. *J. Nat.Prod.* **47**: 373-376.
- [9] **Glddie S., Kellr W.A. and Setterfield G. (1986)**. Production and characterization of somatic hybrids between *Solanum*

- melongena* and *S. sismbriifolium*. *Theor. Appl. genet.* **71**: 613 – 621.
- [10] **Gleddie S., Keller W.A. and Poysa V. (1989)** “Plant regeneration from stem cortex protoplasts of a tomato hybrid”. *Plant Cell Rep.* **8**: 21- 24.
- [11] **Govindan S, Viswanathan S, Vijayasekaran V, Alagappan R (2004)**. Further studies on the clinical efficacy of *Solanum xanthocarpum* and *Solanum trilobatum* in bronchial asthma. *Phytother. Res.* **18**: 805-809.
- [12] **Kao K N and Michayluk M K (1975)** Nutritional requirements for growth of *Vinca hajastana* cells and protoplasts at a very low population density in liquid media; *Planta* **126** 105-110.
- [13] **Keller W.A., Setterfield G., Donglas G., Gleddie S. and Nakamura C. (1992)**.“ Production, characterization and utilization of somatic hybrids of higher plants”. In: Tomes D, Ellis B, Harney P, Kasha M., Petersson R (eds) Application of plant cell and tissue culture to agriculture and industry, University of Guelph, Guelph, ON, pp. 81- 114.
- [14] **Khan TI, Frost S (2001)**. Floral biodiversity: a question of survival in the Indian thar desert. *The Environmentalist*, **21**: 231-236.
- [15] **Kiester D.L. (1975)**. Acetylene reduction by pure cultures of rhizobia. *J Bacteriol.* **123**: 1265- 1268.
- [16] **Kyozuka J, Hayashi Y and Shinomoto K (1987)** High frequency plant regeneration from rice protoplasts by novel nurse culture methods; *Mol. Gen.Genet.* **206** 408-413
- [17] **Larkin PJ, Scowcroft WR (1981)**. Somaclonal variation-a novel source of variability from cell cultures for plant improvement. *Theor. Appl.Genet.* **60**: 197-214.
- [18] **Lee L, Schroll R E, Grimes H D and Hodges T K (1990)** Plant regeneration from indica rice (*Oryza sativa* L.) protoplasts; *Planta* **178** 325-333
- [19] **Maheshwari S C, Gill R, Maheshwari N and Gharyal P K (1986)** The isolation and culture of protoplasts;in *Differentiation of protoplasts and transformed plant cells* (eds) J Reinert and Binding (New York: Springer-Verlag) pp 20-48.
- [20] **Mei-Lei M C T, Reitveld E M, Van Marrewijk G A M and Kool A J (1987)** Regeneration of leaf mesophyll protoplasts of tomato cultivars (*L. esculentum*): factors important for efficient protoplast
- [21] **Murashige T, Skoog F (1962)**. A revised medium for rapid growth and bioassays with tobacco tissue cultures *Physiol. Plant.* **15**: 473-497.
- [22] **Ohnishi T. and Kiyama S. (1987)**. “Increase in yield of mulberry protoplasts by treatment with chemical substance”. *J. Seric. Sci. Japan.* **56**: 407- 410.
- [23] **Ohyama K. and Oka S. (1975)**. “Isolation of protoplasts in *Moraceae*” *Proc. Crop Sci. Soc. Japan.* **44**: 121-122.
- [24] **Sankara Rao K and Gunasekari K (1991)** Control of morphogenesis in tobacco protoplast cultures: Organogenesis Vs. embryogenesis; *Indian J. Biochem. Biophys.* **28** 467-471
- [25] **Schieder O. (1978a)**. “Somatic hybrids of *Datura innoxia* Mill + *Datura discolor* Benth and *Datura innoxia* Mill. + *Datura stramonium* II. Var. *tatula* L.” *Mol. Gen. Genet.*, **162**: 113-119.
- [26] **Shahin E A (1985)** Totipotency of tomato protoplasts; *Theor. Appl. Genet.* **69** 235-240
- [27] **Sinha S, Roy RP, Jha KK (1979)**. Callus formation and shoot bud differentiation in anther culture of *Solanum surattense*. *Can. J. Bot.***57**: 2524-2527.
- [28] **Tewary P.K. and Lakshmisita G. (1992)**. “Protoplasts isolation purification and culture in mulberry (*Morus spp.*)”. *Sericologia*, **32**: 651- 657.
- [29] **Vasil V. and Vasil I.K. (197)**. “Isolation and culture of cereal protoplasts part – 2 embryogenesis and plantlet formation from protoplasts of *Pennisetum americanum*”. *Theor. Appl. Genet.* **56**: 97-99.