

Biotreatment of coloured textile effluents by *Phanerochaete chrysosporium*

P. Jeevana Lakshmi, D.Hymavathi, d Y.Ambethkar

Abstract— Industrial effluents containing dyes pose a threat to human health as they are released into the environment without processing. In the current paper the potential of *Phanerochaete chrysosporium*, a white rot fungus, is investigated for its biological treatment of textile effluents containing three types dyes –Tri phenyl methane, Malachite Green & Reactive Blue. The decrease in optical density values corresponded with decolorization. The effect of pH, temperature, concentrations of cation and anion revealed that the dye decolorization was maximum at pH 6, 28 °C and 0.02 g/Kg concentrations of magnesium sulphate and potassium nitrate.

Index Terms— Decolorization, Dye degradation, Optimization, *Phanerochaete chrysosporium*

I. INTRODUCTION

The usage of dyes has been increased day by day because of increased industrialization and mans' urge for use of various types of colours. Among the 40,000 dyes used in textile industries around 20,000 are accounted to be azo dyes and annual production of these dyes exceeds 7×10^5 tonnes worldwide. Of the total dyes used in the textile industry, 25-50% is lost in the effluents. Presence of dyes in the effluents poses a serious threat to environmental and human health as they are mutagenic and carcinogenic agents. They cannot be completely removed by conventional treatment methods.

However, they can be removed by filtration, coagulation, carbon activated and chemical flocculation methods that generate sludge with secondary disposal problem. Now-a-days, environmental friendly biological processes such as biodegradation, bioaccumulation and bio-sorption for effluent treatment of dyes is receiving increasing interest due to cost effectiveness and ability to produce less sludge. These processes include use of biological material and microorganisms for treatment of dye containing effluents. Bacteria are the widely studied organisms in the microbial degradation of the dyes in the effluents [1]. *Bacillus subtilis* [2] degradation was followed by numerous bacteria such as *Aeromonas hydrophilia* [3], *Bacillus cereus* [4], *Pseudomonas sp.* [5], *E. Coli* [6] etc. However, because of the production of toxic aromatic amines in the anerobic treatment

and dye specific degradation involved aerobic treatment made bacteria less interesting.

This has opened up new arena in search of the other groups of microorganisms involved in dye degradation. In this scenario different groups of fungi have been studied in this decade for their capability of decolorization of dyes from textile effluents. Fungi like *Trametes versicolor* [7], *Trametes hirsute* [8], *Phanerochaete chrysospori* [9], *Schizophyllum commune* and *Lenzites eximia* [10], *Coriolus versicolor* [11], *Trametes versicolor* [12, 13], *Funalia trogii* [14], *Umberopsis isabellina* and *Penicillium geastrivous* [15], *Aspergillus foetidus*, *Rhizopus oryzae* [16] were identified to decolorise polymeric synthetic dyes.

Widely used dyes - Malachite green (MG), reactive blue (RB) and triphenyl methane (TPM) have been studied and their decolorization from the effluents using fungus-*Phanerochaete chrysosporium* has been investigated in the present work. Owing to importance of fungi in decolorization of dye containing effluents, *Phenerochete sps.* isolated from Tirumala Hills, has been studied for the decolorization of the three dye containing effluents- Malachite green, Reactive Blue and Triphenyl methane.

II. MATERIALS AND METHODS

A. Dye decolorization:

Qualitative test of the dye decolorization was done by studying the effect of growth of *Phanerochaete sps.* in the dye incorporated Highley's basal salt medium [17]. 0.5ml of the 48-72 hour old culture isolated on malt yeast extract broth medium was inoculated into the basal media and incubated at 28°C for 8 days in an orbital environmental shaker at 150rpm. The ability of the white rot fungus to decolorize the three types of dyes was visibly recorded.

The flasks were also prepared in a similar way for quantitative measurement of the dye decolorization with all the three types of dyes. After inoculation the flasks were incubated and the samples were read at 600nm for malachite green and at 620nm for reactive blue and triphenyl methane in UV visible spectrophotometer every 48hrs.

B. Optimization of the conditions of decolorization:

Decolorization of textile effluents is effected by various environmental factors like pH, temperature, conc. of cation and anions. To study the effect of pH on the decolorization of the three dyes, each of the effluent flasks were prepared in range of pH 4-9. The flasks are incubated at 28°C in orbital shaking incubator and the readings are taken every 48hrs. Percentage of degradation was calculated as

Manuscript received January 21, 2015.

P. Jeevana Lakshmi, School of Engineering and Technology, Sripadmavati Mahila Visva Vidhyalayam, Tirupati-517502. Andhra Pradesh, India.

D.Hymavathi, Department of Chemical Engineering, S.V. University College of Engineering, Tirupati-517502. Andhra Pradesh, India.

D Y.Ambethkar, Department of Chemical Engineering, Defence Engineering College, Debre Zeit, Oromia, Ethiopia

$$\text{Decolourization\%} = \frac{(C_0 - C_e)}{C_0} \times 100$$

Where, C_0 is initial concentration of dye (mg/l) and C_e is residual dye concentration (mg/l) at different time intervals. To study the effect of temperature on decolorization, each of the effluent flasks were incubated in the temperatures between 25°C and 32°C. The samples were read every 48hrs. To determine the effect of cations and anions, the concentrations of potassium nitrate and magnesium sulphate were adjusted in the range of 0.01 and 0.1 g100ml⁻¹ for the three dye containing textile effluents. The flasks were incubated and the samples were read at 600nm for malachite green and 620nm for reactive blue and triphenyl methane every 48hrs. Percentage decolorization was calculated as given in the earlier section.

III. RESULTS AND DISCUSSION

Phanerochaete sps., white rot fungus, isolated from Tirumala Hills was grown on the three types of dye containing effluents- Malachite green, reactive blue and triphenyl methane. Visible decolorization started from day6 in all the dye containing effluents. Quantitative estimation revealed that optical density decreased (Fig. 1) and percentage of degradation (Fig.2) increased gradually and reached maximum corresponding with observed visual decolorization. Similar decolorization was also seen with the other fungi degrading dyes [18, 19].

Optimization of the conditions of decolorization:

Influence of various environmental factors like pH, temperature, conc. of cations on decolorization of the three dyes by white rot fungus was studied. Effect of pH revealed that the degradation % was optimum at pH 6 and comparatively high degradation is observed at neutral pH (Fig.3). Decolorization of the dye was favoured by slightly acidic pH exhibiting that dye degradation corresponded with nature of the dyes- malachite green, reactive blue and triphenyl methane. Similar degradation pattern was found with the degradation of dyes in acidic pH by other fungal species [20, 21]. The isolate was grown in effluent containing media at different temperatures in the range of 25°C- 35 °C. Optimum decolorization was observed at 28 °C that corresponded with growth of the isolate inferring that dye decolorization is directly affected by the growth and metabolism of the white rot fungus (Fig.4). The effect of concentration of cations and anions on dye decolorization was performed with amendments of concentrations of potassium nitrate and magnesium sulphate in the range of 0.01 to 0.1 g per 100ml in the basal salt medium. It was observed that optimum decolorization was observed with 0.02 concentrations of both the compounds. Any further increase or decrease in the composition of the cations and anions decreased the percentage of decolorization as shown in the Fig.5 and Fig6.




Fig. 1-Optical density values of the dye containing basal media recorded every 24 hour time




Fig.2. % of degradation of the dye decolorization of the three dyes by the *Phenerochaete sps.*




Fig.3. Effect of pH on the decolorization of the three dyes by the *Phenerochaete sps.*




Fig.4. Effect of temperature on the dye decolorization of the three dyes by the *Phenerochaete sps.*




Fig.5. Effect of concentration of KNO₃ on the dye decolorization of the three dyes by the *Phenerochaete sps.*




Fig.6. Effect of concentration of MgSO₄ on the dye decolorization of the three dyes by the *Phenerochaete sps.*

REFERENCES

- [1]. R.V. Kale, M. M. Sable and P. R. Thorat, "Decolorization and degradation of textile azo dye golden yellow HE2R by adapted bacteria", *J. Nat. Env. Poll. Tech.*, Vol. 9(2), 2009, pp.351- 354.
- [2]. H. Horitsu, M.Takada, E. Idaka, M.Tomoyeda & T.Ogawa, "Degradation of p-aminoazobenzene by *Bacillus subtilis*". *Eur. J. Appl. Microbiol.*, Vol. 4, 1977, pp. 217-224.
- [3]. E. Idaka & Y. Ogawa, "Degradation of azo compounds by *Aeromonas hydrophila* var. 2413". *J. Soc. Dyers and Colourists*, Vol. 94, 1978, pp. 91-94.
- [4]. K. Wuhmann, K. I. Mechsner & Th. Kappeler, "Investigation on rate determining factors in the microbial reduction of azo dyes". *Eur. Appl. Microbiol.*, Vol. 9, 1980, pp. 325-338.
- [5]. H. G. Kulla, T. Leisinger, A. W. Cook, R. Hutter & J. Nuesch. "Aerobic bacterial degradation of azo dyes. In *Microbial Degradation of Xenobiotics and Recalcitrant Compounds*", 1981, FEMS Symposium 12, ed. Academic Press, London, pp. 387-399.
- [6]. J. S. Chang, B.Y. Chen, and Y.S. Li, "Stimulation on bacterial decolorization of an azo dye by extracellular metabolites from *Escherichia Col* Strain NO3". *BioresourTechnol.*, Vol. 91(3), 2014, pp. 24 -28.
- [7]. K. Vaidya Varsha and S. Anand Ati, "Decolorization of Reactive Black 5 dye by a white rot fungi *Trametes versicolor*", *J. Nat. Environ. & Pollu. Tech.*, Vol. 8(4), 2009, pp. 701-708.
- [8]. A. Kandelbauer, O. Maute, R.W. Kessler, A. Erlache and M. Gubitz, "Study of dye decolorization immobilized *accase* enzyme-reactor using online spectroscopy", *Biotechnol. Bioeng.*, Vol. 87(4), 2004, pp.52-63.
- [9]. A. Conneely, W. F. Smytha, and G. Mc Mullana, "Metabolism of the phthalocyanine textile dye ramazol turquoise blue by *Phanerochaete chrysosporium*". *FEMS Microbiol. Lett.* Vol. 179, 1999, pp.333-337.
- [10]. K. Selvam, et al., "Biological Treatment Of Pulp And Paper Industry Effluent By White Rot Fungi *Schizophyllum Commune* And *Lenzites Eximia*", *Int. jr. Pharma.Biolog.arch.* Vol.3 (1), 2012, pp. 121-126.
- [11]. S. Sumathi, and B.S. Manju, "Fungal Mediated decolorization of media containing procion dyes", *Water Sci Technol.*, Vol 43(2), 2001. pp. 285-290.
- [12]. Y. Wong, and J. Yu, "Laccase catalysed decolorization of synthetic dyes". *Water Research*, Vol.33 (16), 1999. pp. 3512-3520.
- [13]. J.A. Libra, M. Borchert, and S. Banit, "Competition strategies for the decolorization molasse waste water by *Aspergillus niger*". *Bio resource Technol.*, Vol.57, 2003. pp. 29-235.
- [14]. O. Yesilada, K. Fiskin and E. Yesilada, "The use of white rot fungus *Fungalitrog*(*Mala ty a*) for the decolorization and phenol removal from olive mill waste". *Enviro Technol.*, Vol. 16, 1995, pp.95-100.
- [15]. Q. Yang, M. Yang, K. Pritsch, A. Yediler and A. Kettrup, "Decolorization of synthetic dyes and production of man ganes - dependent peroxidase by new fungal isolate". *Biotechnlgy Letrs* Vol. 25, 2003. pp.709-715.
- [16]. J. K. Polman, and C. R. Breckenridge, "Biomass mediated binding and recovery of textile dyes from waste effluents". *Textil Chemstand colrist.*, Vol. 28 (4), 1996, pp. 31-35.
- [17]. A. Vanden Wymelenberg, et al., "Transcriptome and secretome analysis of *Phanerochaete chrysosporium* reveal complex patterns of gene expression". *Appl. Environ. Microbiol.* Vol. 75, 2009, pp. 4058-4068.
- [18]. S. Senthilkumar, M. Perumalsamy, H. Janardhana Prabhu. "Decolourization potential of white-rot fungus *Phanerochaete chrysosporium* on synthetic dye bath effluent containing Amido black 10B". *Journal of Saudi Chemical Society*.Vol. 18(6), 2014, pp. 845-853.
- [19]. Praphulla Rao et al., "Decolourisation of Synthetic Dyes using *Aspergillus* Species". *Journal of Chemical Engineering and Research*, Vol. 2(1), 2014, pp. 61-68.
- [20]. Amit Kumar Singh, Ritu Singh, Akanksha Soam and Sushil Kumar Shahi "Degradation of textile dye orange 3R by *Aspergillus* strain (MMF3) and their culture Optimization", *International journal current discoveries and innovations*, vol1(1), 2012, pp.7-12.
- [21]. Praveen Sharma, lakhvinder Sharma and Neeraj dilberg, "Biodegradation of orange II dye by *Phanerochaete chrysosporium* in simulated waste water", *Journal of scientific and industrial research*, vol. 6(8), 2009, pp.157-161.