

Long Wave Length tanh Soliton Solution of Sine Gordon Equation

Tapas Kumar Sinha, Joseph Mathew, Sanjib Malla Bujar Baruah

Abstract— Recently H. Sakaguchi and B.A Malomed proposed a novel technique for finding the long wavelength solutions of the Gross Pitaevskii equation. We have applied the technique of Sakaguchi and Malomed to the Sine Gordon equation and derived the equivalent conservation equation. The results are applied to the Josephson junction.

Index Terms— Conservation equation, Josephson junction, Kink solution, Long Wave length soliton solution, Sine Gordon Equation, Soliton, Long Wave length soliton solution

I. INTRODUCTION

Sine Gordon Equation is a partial differential equation which appears in differential geometry and relativistic field theory. The equation, as well as several solution techniques, was known in the 19th century, but the equation grew greatly in importance when it was realized that it led to solutions ("kink" and "antikink") with the collisional properties of solitons. The sine-Gordon equation also appears in a number of other physical applications, including the propagation of fluxons in Josephson junctions (a junction between two superconductors), the motion of rigid pendula attached to a stretched wire, and dislocations in crystals.

the long wavelength region. The effective equation is put in the form of a conservation law (4). The spatial component of the conservation law is an eigenvalue equation. However for Sine Gordon, variable transformations are required to obtain the corresponding Schrodinger equation. The formalism so obtained is applied to the Josephson junction.

II. SINE GORDON EQUATION IN THE LONG WAVE LENGTH LIMIT

The Sine –Gordon equation is

$$\frac{\partial^2 \psi}{\partial t^2} - \frac{\partial^2 \psi}{\partial x^2} + \sin \psi = 0 \quad (1)$$

We look for solutions of the form [1]

Tapas Kumar Sinha, Dept of Computer, NEHU, Shillong , India +919436707015.
 Joseph Mathew, Mathematics, NENU/Union Christian College, Shillong, India,+919436707015.
 Sanjib Malla Bujar Baruah, Mathematics USTM, Khanapara, India +919435387523.

$$\psi(x, t) = \psi^{(0)}(x, t) + \psi^{(1)}(x, t) \cos(2x) + \dots \quad (2)$$

where $\psi^{(0)}(x, t)$ and $\psi^{(1)}(x, t)$ are slowly varying functions of x and t in comparison to $\cos(2x)$ and

$$\psi_{xx}^{(1)}(x, t) \ll \psi^{(1)}(x, t) \quad (3)$$

Using the expansion [8]

$$e^{ix \sin \phi} = \sum_{n=-\infty}^{n=+\infty} J_n(x) e^{in\phi} \quad (4)$$

Where $J_n(x)$ is the Bessel Function

$$\sin \psi = \sin(\psi^{(0)}(x, t) + \psi^{(1)}(x, t) \cos(2x)) \quad (5)$$

$$\cos(x \sin \phi) = \sum_{n=-\infty}^{\infty} J_n(x) \cos n\phi \quad (6)$$

Recently H. Sakaguchi and B. A. Malomed in their seminal paper [1] $\sin(x \sin \phi) = \sum_{n=-\infty}^{\infty} J_n(x) \sin n\phi \quad (7)$

$$\cos(\psi^{(1)}(x, t) \cos(2x)) = \sum_{n=-\infty}^{\infty} J_n(\psi^{(1)}(x, t)) \cos n\left(2x + \frac{\pi}{2}\right) \quad (8)$$

$$\sin(\psi^{(1)}(x, t) \cos(2x)) = \sum_{n=-\infty}^{\infty} J_n(\psi^{(1)}(x, t)) \sin n\left(2x + \frac{\pi}{2}\right) \quad (9)$$

Equating the coefficients of $\cos 2x$

$$\psi_{tt}^{(1)} - \psi_{xx}^{(1)} + 4\psi^{(1)} + \cos(\psi^{(0)}) J_1(\psi^{(1)}) = 0 \quad (10)$$

$$J_1(\psi^{(1)}) = \frac{\psi^{(1)}}{2} - \frac{(\psi^{(1)})^3}{16} \quad (11)$$

Collecting the terms and sub in (11) in (10) one obtain

$$\psi_{tt}^{(1)} - \psi_{xx}^{(1)} + A\psi^{(1)} + B(\psi^{(1)})^3 = 0 \quad (12)$$

Where

$$A = -4.5, B = 0.0625 \quad (13)$$

Note that (13) confirms to the condition for a double well potential well and there by existence of tanh soliton [9].

Let us look for travelling wave solutions of the sine-Gordon equation (5.1) of the form

$$\psi^{(1)} = f(x - vt) \tag{14}$$

Using (14) in (12) we get

$$(v^2 - 1)f'' + Af' + Bf^3 = 0 \tag{15}$$

$$\psi(x, t) = \tanh\left(\frac{x - vt}{\sqrt{2}\zeta}\right) \tag{16}$$

$$\zeta = \frac{1 - v^2}{|A|} \tag{17}$$

Equation (17) represents a localized solitary wave, called kink soliton solution. We note that tanh soliton solution have been found in Josephson's junction (in the long wavelength limit) which are governed by the Sine Gordon equation. Such domain wall soliton solutions have been observed in switching experiments using Nb/Ru/Sr₂RuO₄ junctions [10].

III. CONSERVATION EQUATION

We now derive the conservation equation [2, 4-6] corresponding to (6). Define ρ such that

$$\psi_t^{(1)} = \rho, \psi^{(1)} = \frac{\partial \rho}{\partial x} \tag{18}$$

Using these substitutions in (1) we obtain

$$\rho_t = \rho_{xxx} - A \frac{\partial \rho}{\partial x} + B \left(\frac{\partial \rho}{\partial x}\right)^3 \tag{19}$$

This can be written as a conservation equation

$$\rho_t = \left(\rho_{xx} - A\rho + \frac{B}{4} \left(\frac{\partial \rho}{\partial x}\right)^4 \right)_x = 0 \tag{20}$$

This gives the equation

$$\rho_{xx} - A\rho + \frac{B}{4} \left(\frac{\partial \rho}{\partial x}\right)^4 = C \tag{21}$$

Where c is a constant, We take this c=0

IV. APPLICATION TO JOSEPHSON JUNCTIONS.

Josephson junction consists of two super conductors separated by a thin oxide layer. Each super conductor is characterized by a wave function ψ_1, ψ_2 and phase given by the expressions

$$\psi_1 = \sqrt{n}e^{i\theta_1} \tag{22}$$

$$\psi_2 = \sqrt{n}e^{i\theta_2} \tag{23}$$

Josephson's phase is defined as

$$\psi^{(1)} = \theta_2 - \theta_1 \tag{24}$$

Josephson's phase satisfies the Sine-Gordon equation (1). In the long wavelength limit the Josephson's phase satisfies (5). Josephson's phase is a function of both space and time. The spatial integral defined as

$$\rho = \int \psi^{(1)}(x, t) dx \tag{25}$$

The spatial integral of Josephson's phase is a conserved quantity as derived in (12). Further ρ satisfies the eigenvalue equation (15). We use the boundary condition $\rho \rightarrow 0$ as $x \rightarrow \pm L$. We use the trial solution

$$\rho(x) = Ce^{-kx} \tag{26}$$

$$k^2C - AC - \frac{Bc^4k^4e^{-3kL}}{4} = 0 \tag{27}$$

$$k^2C - AC - \frac{Bc^4k^4e^{3kL}}{4} = 0 \tag{28}$$

Subtracting (26) from (25) we obtain

$$\frac{Bc^4k^4}{2} \sinh(3kL) = 0 \tag{29}$$

This implies $3kL = -in\pi$ (30)

Since L is real k must be imaginary. In other words (25) represents an oscillatory solution. Let

$$k = -i\kappa \tag{31}$$

Where κ is a real quantity. Substituting (30) in (29) we get

$$\kappa = \frac{n\pi}{3L} \tag{32}$$

This defines wave-vectors in (26). Note that only certain wave vectors are permissible. This is expected, as we are in the quantum domain where only discrete states are allowed.

V. CONCLUSION

We found the long wave length kink soliton solution of Sine Gordon equation. We have derived the conservation equation of Sine Gordon equation in the long wavelength limit. The kink solution in this limit as well as conservation laws in this limit are derived. The results are applied to the Josephson's junction and we found Josephson's phase exhibits spatial sinusoidal oscillations. Further we find that the spatial integral of the Josephson's phase is a conserved quantity.

REFERENCES

[1] Hidetsugu Sakaguchi and Boris A. Malomed, Phys. Rev. E 72,046610 -1(2005).
 [2] T.C. Au-Yeung, C.Au, and P.C. W, Fung Phys. Rev. 29, 2370(1984).
 [3] Thornsten Holst and J. Bindslev Hansen, Physical Review B, 44 2238-2245 (1991).

- [4] C.S. Gardner, J.M. Greene, M.D. Kruskal, and R.M. Miura, Method for Solving the Korteweg-deVries Equation, Phys. Rev. Lett.19(1967), 1095-1097.
- [5] R.M. Miura, Korteweg-deVries Equation and Generalizations. I. A Remarkable Explicit Non-linear Transformation, J. Math. Phys.9 (1968), 1202-1204.
- [6] P.D. Lax, Integrals of Nonlinear Equations of Evolution and Solitary Waves, Comm. Pure Appl. Math. 21 (1968), 467-490.
- [7] Available in: en.wikipedia.org/wiki/Sine-Gordon_equation
- [8] Available at "http://en.wikipedia.org/wiki/Bessel_function
- [9] J. A Krumhansl and J.R Schrieffer Physical Review B Vol.11 No 9, 1 May 1975
- [10] Available at:
www.nature.com/srep/2013/130821/srep02480/full/srep02480.html

Tapas Kumar Sinha, the second author obtained his M.Sc(Physics) BHU, MS(Physics) and MS(BioPhysics) from University of Cincinnati, USA PhD in Neural Networks in 1995 from North Eastern Hill University, Shillong, India. Currently he is Associate Professor (Computer Science) at North Eastern Hill University. His major field of interest is applications of Nonlinear Dynamics. Currently he is working on applications of Nonlinear Dynamics to refractive photonic media, development of nonlinear optical filters and massively parallel optical computers using such media. Dr. Sinha has worked extensively as a Software Consultant for most major companies (IBM, Mead Paper, Reynolds Tobacco, and ITT) in the U.S. Currently he is guiding three students for Ph.D.

Joseph Mathew the co-author is a post graduate in Mathematics (M.Sc.) and Computer Science (MCA) working in Union Christian College, Shillong as Associate Professor in the department of Mathematics and Vice Principal. His major field of interest is applications of Nonlinear Dynamics. Currently he is working on applications of Nonlinear Dynamics to refractive photonic media, development of nonlinear optical filters and massively parallel optical computers using such media. He is currently working under Dr. Sinha and Dr. Sanjib Malla Bujar Baruah for his PhD.

Sanjib Malla Bujar Baruah, is the Head of the department of Mathematics, in the University of Sc. and Technology, Khanapara, Meghalaya India. He has completed his PhD from Dibrugarh University. He is guiding Joseph Mathew for his PhD in Mathematics at USTM.